DSIP

Q. P. Code: 12213

(Time: 2½ Hours)

[Total Marks: 75]

N.B.		l questions are compulsory.					
	2) Figures to the right indicate marks.						
	3) Illi	ustrations, in-depth answers and diagrams will be appreciated.					
	4) Mi	xing of sub-questions is not allowed.	\$ 8 C				
	5) Us	se of own non-programmable calculator is allowed.					
Q. 1	Atte	Attempt All(Each of 5Marks)					
(a)	Multiple choice questions						
	i. The range of correlation coefficient is						
		A. 0 to 1 B1 to 1 C1 to 0 D. None of the above					
	ii.	If B is subset of A then $p(A/B) = \dots$	352				
		A. 1 B. P(A) C. P(B) D. None of the above					
	iii.	In less than type ogive curve, the points are plotted for	E.				
	12	A. the lower boundary and frequency.					
		B. the upper boundary and cumulative frequency.					
	8.5	C. the lower boundary and cumulative frequency.					
	1	D. None of the above					
	iv.	The measure of central tendency which can be used for further mathematical treatment is					
		B. Median					
	4	C. Mode					
	0	D. All the above					
	0.4						
		If the lower and upper limits of the class interval are 20 and 30					
3	C. C.	respectively then the class mark will be					
SAL		A. 10 D. 25					
(b)							
ינטן		the blanks					
20.0		Mode is the occurring value in data set.					
300	0 66	In histogram the width of the bar will be decided on the basis of					
100	iii.	If the correlation coefficient between two variables X and Y is					
970	3 4	perfect then the correlation coefficient r =					
5 4 0	iv	For Y = a + bx, Y is called asvariable					
		$p(A\cap A') = \dots$					

Short answers.

- Write two requisites of good measure of central tendency.
- Define variance. ii.
- Write the formula of regression coefficient of X on Y.
- Define probability.
- Define mutually exclusive events

Q. P. Code: 12213

Attempt the following (Any THREE)(Each of 5Marks) Q. 2 (15M)Explain with one example Nominal scale, Ordinal scale and ratio scale. (a) (b) Write a short note on Frequency polygon.ii. Stem and leaf plot. Given the following data on the marks obtained by students in some (c) examination. 22,24,15,25,10,12,14,8, 2, 4, 4,6,12,14,16,17,18,18,17,14,10,10,8,9,22,21,23,20,18. Construct frequency distribution with inclusive type class interval. Obtain mean and mode for the following data (d) C.I. 10-20 20-30 30-40 40-50 50-60 60-70 Frequency Explain the procedure of obtaining Quartile deviation for grouped data. (e) Define standard deviation and Find coefficient of variation for the (f) following dataX: 12,13,14,15,16,12,14,16,13,15,14,14,12 Q. 3 Attempt the following (Any THREE) (Each of 5Marks) (15M)Define first four raw moments about origin zero and central moments of a (a) distribution. Also state the relationship between raw and central moments Explain the concept of skewness. Also distinguish between positive and (b) Negative skewness. With usual notation μ_1 =2, μ_2 =8, μ_3 =14 and μ_4 =50 then Compute β_1 and (c) Represent Positive, Negative and Perfect correlation using scatter plots. (d) Explain the concept of correlation and regression. Also comment, how (e) regression is different from correlation. (f) For the following data obtain the regression line of the type Y on X 12 14 16 15 18 3 Attempt the following (Any THREE) (Each of 5Marks)

(15M)

(a) Define the Following with one example:-

i. Random Experiment with one example.

ii. Sample space and Event with one example.

- A ticket is drawn from a box containing 30 tickets and a number on it is observed. Obtain the probability that ticket drawn has a number
 - Less than 6
 - Greater than 20 ii.
 - iii. Multiple of 5.
- The letter of the word 'EQUATION' are arranged randomly. What is the probability that an arrangement
 - i. Starts and ends with vowel.
 - ii. Have all vowels together.
 - iii. State Addition theorem and Bay's theorem.
- State Addition Theorem and Bay's Theorem.

Q. P. Code: 12213

- (e) Two dice are thrown simultaneously. Find the probability that the sum being 6 or same number on both dice.
- (f) A hospital has 3 doctors X,Y&Z operating independently. The probability that doctor X is available is 0.9 and that for Y is 0.6 and for Z is 0.7; What is the probability that at least one doctor is available when needed?
- Q. 5 Attempt the following (Any THREE) (Each of 5Marks)

(15M)

- (a) Explain the procedure of plotting Bar chart and Pie Chart.
- (b) Write two merits and two demerits of the Mode and Coefficient of range.
- (c) Define Kurtosis and explain different types of kurtosis.
- (d) Obtain Spearman's Rank correlation between performance in Maths and Computer Science. The scores are given below:

Computer science.	Tie bee	0	59 N. Car Car	1	17.5	P 32 3 C
Maths	56	65 72	48	56	70	68
Computer Science	76	60 50	75	66	87	77

- (e) If the two regression equations are 4y 5x 33 = 0 and 20y 9x 107 = 0, Find: i. Mean of x and y ii. Correlation coefficient between x and y.
- (f) Stockiest has 20 items in a lot. Out of which 12 are non-defective and 8 are defective. A customer selects 3 items from the lot. What is the probability that out of these three items:
 - i. Three items are non-defective
 - ii. Two are non-defective and one is defective
